



Review

# A Systematic Review on the Use of Wearable Body Sensors for Health Monitoring: A Qualitative Synthesis

Annica Kristoffersson \* D and Maria Lindén

School of Innovation, Design and Engineering, Mälardalen University, 722 20 Västerås, Sweden; maria.linden@mdh.se

\* Correspondence: annica.kristoffersson@mdh.se

Received: 4 February 2020; Accepted: 5 March 2020; Published: 9 March 2020



Abstract: The use of wearable body sensors for health monitoring is a quickly growing field with the potential of offering a reliable means for clinical and remote health management. This includes both real-time monitoring and health trend monitoring with the aim to detect/predict health deterioration and also to act as a prevention tool. The aim of this systematic review was to provide a qualitative synthesis of studies using wearable body sensors for health monitoring. The synthesis and analysis have pointed out a number of shortcomings in prior research. Major shortcomings are demonstrated by the majority of the studies adopting an observational research design, too small sample sizes, poorly presented, and/or non-representative participant demographics (i.e., age, gender, patient/healthy). These aspects need to be considered in future research work.

**Keywords:** health monitoring; IoT; physical activity monitoring; qualitative synthesis; remote health management; research shortcomings; sensor systems; user demography; wearable body sensors

#### 1. Introduction

The use of wearable body sensors for health monitoring as a means for supporting clinical and remote health monitoring in real-time and to provide health trend monitoring with the aim to predict/prevent health deterioration has the potential to lower the burden on the healthcare system and thereby reduce healthcare costs. The number of available wearable and wireless body sensors and systems are rapidly growing. Simultaneously, research on more energy-efficient and more accessible/smaller sensors for acquiring data as well as research on automatic data analysis of the Big Data, which the sensor-based systems are expected to generate, is being conducted. This advanced data analysis has the potential of generating personalized diagnoses and providing recommendations on treatments at a personalized level. While a promising area, we argue that the data collected for generating advanced data analysis algorithms need to come from participants representing the expected users of these systems.

This systematic review provides a qualitative synthesis of the articles retrieved on using wearable body sensors for health monitoring. We analyze the articles from many perspectives including author affiliations in countries, publication years, context of use, sensor category, research methodology, sample sizes, and participant demographics (i.e., age, gender, patient/healthy). This analysis has identified a number of shortcomings in prior research with respect to both sample size, but also to participant demographics where the latter strongly affects the validity of the results. These shortcomings need to be considered in future research, not only for understanding the user experience, but also to ensure that the advanced data analysis algorithms can reason on data which are representative and valid for the expected users of the systems.

Sensors **2020**, 20, 1502 2 of 21

#### 2. Methodology

Following the requirements of MDPI Sensors, a systematic review following the PRISMA guidelines [1] was conducted. A total of seven databases were searched, including: Web of Science Core Collection, MEDLINE, Scopus, ScienceDirect, Academic Search Elite, ACM Digital Library, and IEEE Xplore.

The searches were conducted on 24–25 April 2019. The search phrases resulting in the identification, and addition to an EndNote database, of related articles are shown in Table 1. During the search, the keywords were changed in order to broaden or narrow the number of articles found using the previous search phrase. For example, "Ecare" or "mHealth" or "ehealth" was replaced with "care" or "Health" in the second search in Web of Science Core Collection. The same search phrase was used for MEDLINE but it resulted in thousands of hits in SCOPUS. Several additional searches aiming at limiting the number of hits were conducted resulting in "care" or "Health" being replaced with the original search phrase "ecare or mhealth or ehealth" and the exclusion of "feedback" and "pilot application". The search phrase used for Scopus resulted in no hits in Science Direct. Therefore, two less narrow searches were conducted. Variations of these phrases were used in Academic Search Elite, ACM Digital Library and IEEE Xplore.

**Table 1.** An overview of search phrases and databases used during article retrieval.

Database	Search Phrase	Number of Articles
Web of Science Core Collection	ALL FIELDS: (("body sensor" or "wireless body sensor" or "wireless wearable technology" or "biomedical sensor" or "IoT") and ("Ecare" or "mHealth" or "eHealth") and ("Social impact" or "Compliance" or "Acceptance" or "Clinical trial" or 'Pilot test" or 'Human input" or "Feedback" or "Pilot application" or "Human in the loop"))	7
Web of Science Core Collection	ALL FIELDS:(("body sensor" or "wireless body sensor" or "wireless wearable technology" or "biomedical sensor" or "IoT") and ("care" or "Health") and ("Social impact" or "Compliance" or "Acceptance" or "Clinical trial" or "Pilot test" or "Human input" or "Feedback" or "Pilot application" or "Human in the loop"))	142
MEDLINE (Web of Science)	TOPIC: (((((("body sensor") OR "wireless body sensor") OR "wireless wearable technology") OR "biomedical sensor") OR "IoT") AND ("care") OR "Health")) AND ((((((("Social impact") OR "Compliance") OR "Acceptance") OR "Clinical trial") OR "Pilot test") OR "Human input") OR "Feedback") OR "Pilot application") OR "Human in the loop")) Timespan: All years. Indexes: MEDLINE.	25
Scopus	ALL(body sensor OR wireless body sensor OR wireless wearable technology OR biomedical sensor) AND (ecare OR mhealth OR ehealth) AND (Social impact OR compliance OR acceptance OR Clinical trial OR Pilot test) Limiting to English	187
ScienceDirect	Title, abstract, keywords: "wearable sensors" and health and impact. Limited to review articles, research articles, conference abstracts, case reports.	13

Sensors 2020, 20, 1502 3 of 21

Table 1. Cont.

Database	Search Phrase	Number of Articles
ScienceDirect	Title, abstract, keywords: "body sensor" and health and impact. Limited to review articles, research articles, conference abstracts, case reports.	5
Academic Search Elite	Free text search: "body sensor" and health and impact English.	8
Academic Search Elite	Free text search: "body sensor" and health and acceptance	3
<b>ACM</b> Digital Library	(+"body sensor" +and +health +and +impact)	12
IEEE Xplore	"body sensor" and health and impact	81
IEEE Xplore	"body sensor" and health and trial	12

Article Selection, Inclusion and Exclusion Criteria

The search resulted in 495 articles. Thereafter, the articles were screened in several steps using EndNote:. Thirty duplicated articles were eliminated and 288 articles were excluded after reviewing each title and abstract individually. Abstracts and articles retrieved that did not match the main research question were excluded from further consideration. For example, we excluded articles on studies using solely environmental exposure sensors or smart home sensors.

Then, pdf copies of all remaining articles were downloaded. Copies of abstracts, introductions and conclusions were extracted to OneNote after which an additional screening was conducted. The eligibility criteria for inclusion in the review were:

- Articles should be published as a journal article or in conference proceedings.
- Articles should consider wearable technology and monitoring.
- Articles should present results from studies where sensor data were collected using humans.
   Alternatively, the articles present information on a system where the user trial is planned for but not conducted yet.
- Articles should be in English.

Overviewing the remaining 177 articles, it was found that the number of publications relating to some health conditions, henceforth called article categories, was low. Therefore, no articles were excluded based on publication year. In addition, we excluded the numerous review articles from further analysis as they cannot be considered original research, i.e., the review articles retrieved were excluded since they do not directly report on a conducted study of people or on the planning of such a study. Publications that met the inclusion criteria, and therefore, considered for further reviewing were 73. The study selection process is depicted in Figure 1.

Sensors 2020, 20, 1502 4 of 21




Figure 1. The article selection process.

## 3. Qualitative Synthesis

Inspired by Kekade et al.'s review from 2018 [2], we conducted a qualitative synthesis of the 73 included research articles. They were published between 2010 and 2019, i.e., spanning approx. 9.5 years, among which one article was published in 2010, two in 2011, seven in 2012, two in 2013, seven in 2014, twelve in 2015, nine in 2016, fourteen in 2017, fourteen in 2018 and five before April 24th 2019, see Figure 2. In average, 7.6 articles were published per year during the period 2010–2018. The authors of the 73 research articles were affiliated in 32 countries representing six continents (Africa, Asia, Australia, Europe, North America and South America). See Figure 3 and 4 for further information on which countries authors are affiliated in and the number of publications per country with affiliated authors. The articles were sorted into the following article categories: Asthma/COPD, Cardiovascular diseases, Diabetes and nutrition, Gait and fall, Neurological diseases,

Sensors **2020**, 20, 1502 5 of 21

Physical activity recognition, Rehabilitation, and Stress and sleep. All articles not directly related to any of the aforementioned article categories were sorted into an article category named Additional. Figure 5 depicts the category-wise distribution of the selected articles. Studying the distribution of articles related to health and physical activity monitoring respectively, it can be seen that 47 % of the articles were related to health (Asthma/COPD, Cardiovascular diseases, Diabetes and Nutrition, Neurological diseases, and Stress and sleep). As much as 39 % of the articles were related to physical activity monitoring (Gait and fall, Physical activity recognition, and Rehabilitation). It is unclear why such a large portion of the articles were related to physical activity monitoring. Possible reasons include that it is easier to monitor physical activity using sensors whereas measures relating to health, e.g., vital signs, need to be provided in a more timely manner.






Figure 2. Number of articles per year. \* only the articles published prior to 24 April 2019 are counted.

Sixty research articles reported on studies conducted with people at some level, these are reported in Table 2. We categorized the sensors according to the sensor categories used in [2], namely, physical activity, vital signs, electrocardiography (ECG) and other. Studies reporting on devices measuring movement or activity were classified under the sensor category physical activity. Vital signs include the parameters: blood pressure (BP), body temperature (BT), respiratory rate (RR), heart rate (HR)/pulse, and peripheral oxygen saturation (SpO<sub>2</sub>). Studies measuring ECG were classified under ECG. Finally, studies using sensors for diabetes, swallowing, etc., or a combination of sensors from several sensor categories were classified under the sensor category other. The sensor categories physical activity and other include 23 studies each, vital signs includes three studies, and ECG includes ten studies reported upon in seven research articles.

Similarly to Kekade et al. 2018 [2], we also assessed the studies' reporting of research design (Table 2), and the reported participant demography, i.e., number of participants, age, gender and the distribution of healthy participants and patients (see Sections 3.1–3.4 and Table 3). Many studies presented the participant demographics poorly, or not at all [3–12]. Rather than excluding these from the tables, we indicate missing information with a "-". However, we question the fact that all these studies were accepted for publication without providing any information on the participants. Our findings are further discussed in Section 4.



**Figure 3.** Number of authors affiliated in each country. Authors are calculated for each article, i.e., an author may be calculated more than once and in more than one country.



**Figure 4.** Number of articles per country. Papers with several authors may be counted for several countries.

**Table 2.** List of articles reporting on conducted studies. —indicates that information is missing.

Author, Year	Ref.	<b>Article Category</b>	Research Design	No. of Participants	Sensor Category	
Bonnevie et al. 2019	[13]	Asthma/COPD	Observational	104	Vital signs	
				5		
Caulfield et al. 2014	[14]	Asthma/COPD	Observational	10	Physical activity	
Estrada et al. 2016	[15]	Asthma/COPD	Observational	1	Other	
Katsaras et al. 2011	[16]	Asthma/COPD	Randomized control	48	Other	
Naranjo-Hernández et al. 2018	[17]	Asthma/COPD	Observational	2	Vital signs	
				9		
Huang et al. 2014a	[18]	Cardiovascular diseases	-	225	ECG	
Huang et al. 2014b	[19]	Cardiovascular diseases	Case-control	84	ECG	
Javaid et al. 2018	[20]	Cardiovascular diseases	Observational	60	Other	
Li et al. 2019	[3]	Cardiovascular diseases	Observational	16	Other	
Raad et al. 2015	[21]	Cardiovascular diseases	-	30	ECG	
			-	2		
Simjanoska et al. 2018	[22]	Cardiovascular diseases	Observational	16	ECG	
,				3		
				25		
				7	Dataset ECG	
Susič and Stanič 2016	[23]	Cardiovascular diseases	-	13	ECG	
Al-Taee et al. 2015	[24]	Diabetes and nutrition	-	22	Other	
Alshurafa et al. 2014 and Alshurafa et al. 2015	[25,26]	Diabetes and nutrition	Observational	10	Other	
				20		
Dong and Biswas 2017	[27]	Diabetes and nutrition	Observational	14	Other	
Onoue et al. 2017	[28]	Diabetes and nutrition	Randomized control	101	Physical activity	
Atallah 2012	[29]	Gait and fall	Observational	34	Physical activity	
Godfrey et al. 2014	[30]	Gait and fall	Observational	24	Physical activity	
Lee et al. 2015	[31]	Gait and fall	Observational	11	Physical activity	
Liang et al. 2012	[32]	Gait and fall	Observational	8	Physical activity	
Liang et al. 2018	[33]	Gait and fall	Observational	18	Physical activity	
Paiman et al. 2016	[34]	Gait and fall	Observational	2	Other	
Tino et al. 2011	[35]	Gait and fall	Observational	3	Other	
Williams et al. 2015	[36]	Gait and fall	Observational	5–6	Physical activity	
Wu et al. 2013	[4]	Gait and fall	Observational	7	Physical activity	
Wu et al. 2019	[37]	Gait and fall	Observational	15	Physical activity	
Zhao et al. 2012	[38]	Gait and fall	Observational	8	Physical activity	
Zhong et al. 2019	[39]	Gait and fall	Observational	56	Physical activity	

 Table 2. Cont.

Author, Year	Ref.	Article Category	Research Design	No. of Participants	Sensor Category	
Giuberti et al. 2015	[40]	Neurological diseases	Observational	24	Physical activity	
Gong et al. 2015, Gong et al. 2016	[41,42]	Neurological diseases	Case-control	41	Physical activity	
Kuusik et al. 2018	[43]	Neurological diseases	Observational	51	Physical activity	
Sok et al. 2018	[44]	Neurological diseases	Observational	13	Physical activity	
Stamate et al. 2017 and Stamate et al. 2018	[45,46]	Neurological diseases	Observational	12	Other	
Castro et al. 2017 and Rodriguez et al. 2017	[5,6]	Physical activity recognition	Observational	3	Other	
Doron et al. 2013	[7]	Physical activity recognition	Observational	65	Other	
		, , ,		20		
Rednic et al. 2012	[47]	Physical activity recognition	Observational	17	Physical activity	
Xu et al. 2014	[8]	Physical activity recognition	Observational	14	Other	
Xu et al. 2016	[9]	Physical activity recognition	Observational	4	Other	
		, , ,		3	Physical activity	
				5	,	
				6		
Argent et al. 2019	[48]	Rehabilitation	Observational	15	Physical activity	
Banos et al. 2015	[49]	Rehabilitation	Observational	10	Other	
Lee et al. 2018	[50]	Rehabilitation	Case-control	30	Physical activity	
Timmermans et al. 2010	[51]	Rehabilitation	Observational	9	Physical activity	
Whelan et al. 2017	[52]	Rehabilitation	Observational	55	Physical activity	
Xu et al. 2017	[53]	Rehabilitation	Observational	6	Other	
Lin et al. 2012	[54]	Stress and sleep	Case-control	18 (6/12)	Physical activity	
Nakamura et al. 2017	[55]	Stress and sleep	Observational	4	Other	
Parnandi and Gutierrez-Osuna 2017	[56]	Stress and sleep	Randomized control	25	Other	
Uday et al. 2018	[57]	Stress and sleep	Observational	10	Other	
Umemura et al. 2017	[58]	Stress and sleep	Case-control	54	Other	
Velicu et al. 2016	[10]	Stress and sleep	Observational	-	-	
Ayzenberg and Picard 2014	[59]	Additional	Crossover	10	Other	
Pagán et al. 2016	[60]	Additional	Observational	2	Other	
Rawasdeh et al. 2017	[61]	Additional	Observational	55	ECG	
Seeger et al. 2012	[11]	Additional	-	-	Other	
Wannenburg and Malekian 2015	[12]	Additional	Observational	4–8	Vital signs	
Wu et al. 2018	[62]	Additional	Observational	20	ECG	


**Table 3.** Demographic information on conducted studies. - indicates that information is missing.

Ref.	Article Category	No. of Participants	Age Group	Age Statistics	Male	Female	Patient	Healthy
[13]	Asthma/COPD	104	57–70	64	67 (64%)	37 (36%)	104	
		5	50-66	62	- ` ′	- ` ´	5	
[14]	Asthma/COPD	10		$61.5 \pm 5.7$	5	5	10	
[15]	Asthma/COPD	1	-	-	1			1
[16]	Asthma/COPD	48	-	-	48		48	
[17]	Asthma/COPD	2	36 and 42		2			2
		9	55–76	$64 \pm 6.6$	6	3	9	
[18]	Cardiovascular diseases	225	-	-	-	_	225	
[19]	Cardiovascular diseases	84	_	-	_	_	1 group	1 group
[20]	Cardiovascular diseases	60	_	$26.9 \pm 6.1$	28	32	0 1	60
[3]	Cardiovascular diseases	16	-	-	-	_	-	-
[21]	Cardiovascular diseases	30	20-23		-	_	-	-
		2	_	-	_	_	2	
[22]	Cardiovascular diseases	16	16–72	-	_	_	-	_
		3	25-27	-	-	_	-	-
		25	20-73	-	_	_	14	11
		7	20-74	-	_	_		7
[23]	Cardiovascular diseases	13	_	$50.6 \pm 9$	8	5		13
[24]	Diabetes and nutrition	22	-	-	-	_	22	
[25,26]	Diabetes and nutrition	10	20-40		8	2	-	_
. , .		20	20-40		12	8	-	_
[27]	Diabetes and nutrition	14	_	-	9	5		14
[28]	Diabetes and nutrition	101	_	$57.1 \pm 12.5$	56	45	101	
[29]	Gait and fall	34	_	$28.22 \pm 12.77$	21	13		34
[30]	Gait and fall	24 (12/12)	20-40	$32.5 \pm 4.8$	7	5		12
		, ,		$65.0 \pm 8.8$	5	7		12
[31]	Gait and fall	11	_	$27.6 \pm 4.3$	11			11
[32]	Gait and fall	8	-	$23 \pm 3.45$	8			8
[33]	Gait and fall	18	_	$25 \pm 3.24$	12	6		18
[34]	Gait and fall	2	28 and 24	-	1	1		2
[35]	Gait and fall	3	40-70	-	_	_	-	_
[36]	Gait and fall	5-6 (1/5)	27	-	1		-	_
		· /	21–36	27	4	1	-	-
[4]	Gait and fall	7	-	-	-	-	-	-
[37]	Gait and fall	15	20-27	-	-	-		15
[38]	Gait and fall	8	_	$28.5 \pm 4.3$	-	-		8
[39]	Gait and fall	56 (28/28)	-	$24.6 \pm 2.7$	14	14		28
		,	>55	$66.1 \pm 5.0$	18	10		28

 Table 3. Cont.

Ref.	Article Category	No. of Participants	Age Group	Age Statistics	Male	Female	Patient	Healthy
[40]	Neurological diseases	24	31–79	$65.9 \pm 12.3$	17	7	24	
[41,42]	Neurological diseases	41 (28/13)	-	$40.5~\pm~9.4$	25%	25%	28	13
			-	$39.3 \pm 10.3$	47%	53%		
[43]	Neurological diseases	51	-	-	-	-	51	
[44]	Neurological diseases	13	22-50	-	9	4	13	
[45,46]	Neurological diseases	12	-	-	-	-	12	
[5,6]	Physical activity recognition	3	-	-	-	-	-	-
[7]	Physical activity recognition	65	-	-	-	-	-	-
	, , ,	20	-	-	-	-	-	-
[47]	Physical activity recognition	17	-	-	10	7	-	-
[8]	Physical activity recognition	14	-	-	-	-	-	-
[9]	Physical activity recognition	4	-	-	-	-	-	-
	, , , ,	3	-	-	-	-		3
		5	-	-	-	-	5	
		6	-	-	3	3	-	_
[48]	Rehabilitation	15	-	$63 \pm 8.32$	6	9	15	
[49]	Rehabilitation	10	21-37	-	8	2	_	_
[50]	Rehabilitation	20	-	$54.4 \pm 10.1$	-	-	20	
		10		$53.8 \pm 11.4$	-	-		10
[51]	Rehabilitation	9	-	60.7	5	4	9	
[52]	Rehabilitation	55	_	$24.21 \pm 5.25$	37	18		55
[53]	Rehabilitation	6	_	$72.5 \pm 6.0$	3	3	_	_
[54]	Stress and sleep	18 (6/12)	19–22 overall	-	5	1	_	_
	1	( , ,			11	1	_	_
[55]	Stress and sleep	4	25–36	-	4			4
[56]	Stress and sleep	25	19–33	-	15	10	_	_
[57]	Stress and sleep	10	_	-	-	_		10
[58]	Stress and sleep	54 (26/28)	-	22	-	-		54
. ,	1	, ,	_	21	-	_	_	_
[10]	Stress and sleep	-	_	-	_	_	_	_
[59]	Additional	10	25–35	$30.8 \pm 4.2$	9	1		10
[60]	Additional	2	-	_		2	2	
[61]	Additional		18–22	-	50%	50%	-	_
[11]	Additional	-	_	-	_	_	_	_
[12]	Additional	4-8 (4/4)	-	-	_	_	_	_
r1		- (-/ -/	-	-	_	_	_	_
[62]	Additional	20	-	-	_	_		20

Sensors 2020, 20, 1502 11 of 21



**Figure 5.** Category-wise distribution of the selected articles. Number of articles for Additional = 10, Asthma/COPD = 6, Cardiovascular diseases = 8, Diabetes and nutrition = 5, Gait and fall = 15, Neurological diseases = 8, Physical activity recognition = 7, Rehabilitation = 7, Stress and sleep = 7.

For completeness, the remaining 13 articles not listed in Table 2 and 3 were distributed over eight article categories: Asthma/COPD [63], Cardiovascular diseases [64], Gait and fall [65–67], Neurological diseases [68], Physical activity recognition [69], Rehabilitation [70], Stress and sleep [71], and Additional [72–75]. Six articles report on systems where studies are upcoming [63,64,72–75]. One of them [64] is a continuation of the study reported in [23]. Three articles report on studies using datasets [66,67,69]. Two articles report on qualitative studies of observational and/or interview nature [68,70]. The continuation of the qualitative study [70] is reported upon in [48]. The evaluation in [65] is not clearly presented and the system developed in [71] uses wearable body sensors only to collect ground truth data for a contactless sleep monitoring system. Therefore, [71] was excluded from further qualitative analysis.


#### 3.1. Research Methodology

Table 2 reports on the four research designs identified while analyzing the research articles: case-control, crossover, randomized control and observational. Articles categorized as adopting a case-control research design are prospective and include studies with two groups. In most articles, one group is a healthy control group and the other a group sharing an illness. However, in this review, also articles comparing the measures for two distinct groups (e.g., non-shift workers in rural and urban areas) have been categorized as adopting a case-control research design. Articles categorized as adopting a randomized-control research design have participants with the same background being randomly assigned to one of two study conditions. One article has been categorized as a crossover study [59], the participants have experienced both study conditions but in randomized order. The articles categorized as being observational are typically conducted in a controlled fashion during which data are collected. In this review, the majority of the articles were categorized as being observational. A few articles adopted a case-control [19,41,42,50,54] or randomized control research design [16,28,56]. For some articles [11,18,21,23,24,58], information provided on how the experiments were conducted was not sufficient for determining the research design adopted.

Studying the number of participants included in the studies, we first summarized the number of participants in the cases where an article reported on several smaller studies. It can be seen from Figure 6 that 57% of the studies were conducted with up to 20 participants and that 30% were conducted with 10 or fewer participants. Only 40% of the studies were conducted with 21 or more participants (22% collected 21-50 participants, 13% had 51–100 participants leaving 5% with more than 100 participants).


Looking more closely into each article category, Figure 7 shows that the majority of the studies within the categories Asthma/COPD, Gait and fall, Physical activity recognition, Rehabilitation, Stress and sleep, and Additional were conducted with up to 20 participants. The studies with more than 100 participants fall within the categories Asthma/COPD, Cardiovascular diseases, and Diabetes and nutrition. Studies with 51–100 participants were conducted within the categories Cardiovascular diseases, Gait and fall, Neurological diseases, Rehabilitation, Stress and sleep, and Additional.

To make technical validations that a sensor is working, a small number of participants can be accepted. However, to be used in clinical investigations, power calculations taking the research question into account should be used to decide the number of needed participants.



**Figure 6.** Distribution of the number of participants per included study. - denotes studies which did not provide information on number of participants.

Sensors 2020, 20, 1502 13 of 21



**Figure 7.** Distribution of the number of participants per article category. - denotes studies which did not provide information on number of participants.

### 3.2. Age Distribution

Information on the participants' age was provided in 35/60 (58.3%) of the articles reporting on data collection studies with people (Table 3). Another two articles [21,23] provided the information on age for only one of the study groups. A very limited number of studies were conducted with people where  $\mu_{age} > 65$  [40,53] or  $\mu_{age} > 60$  [13,14,48,51]. Two studies [30,39] were conducted with one young group and one group where  $\mu_{age} > 65$ , whereas  $\mu_{age} > 60$  for one of the groups in [17]. Two articles report on studies with large age ranges where some participants exceed 65 years of age (16–72 and 20–73 in [22], and 40–70 in [35]).

Studying the articles from an article category perspective, none of the studies reporting on the categories Cardiovascular diseases, Diabetes and nutrition, Other or Stress and sleep was conducted with participants where  $\mu_{age} > 60$ . The categories Asthma/COPD, Gait and fall, Neurological diseases, and Rehabilitation include some studies with this age group. None of the studies within the Physical activity recognition category report on the participants' age.

#### 3.3. Gender Distribution

Information on the participants' gender was provided in 33/60 (55%) of the articles reporting on data collection studies with people (Table 3). Three more articles [9,13,23] reported on studies with more than one group but not the gender for all groups.

Studying the articles from an article category perspective, all Asthma/COPD studies except [13] provided full information on gender distribution. The latter, [13] also reports on a study with a subset of the participants without providing information on gender. Regarding cardiovascular diseases studies, only [20,23] provided full information on gender distribution. Another 20 want to participate in screening although the study described in [23] is not approved yet by an ethical committee. All but one study within Diabetes and nutrition report on gender. The majority of the studies within Gait and fall contain information on gender. More than half (57%) of the articles on Neurological diseases and 50% of the articles on Other present information on gender. Regarding the category Physical activity recognition, only one article [47] provides full information on gender. Another article, [9] provides information on gender for one of their four sub-studies. The majority (80%) of the Rehabilitation studies and 50% of the Stress and Sleep studies provide gender information.

Studying the articles from a gender distribution perspective, the vast majority of the participants in the studies reporting on Asthma/COPD are men. For Cardiovascular diseases, [20] had a rather

even gender distribution, [23] reported on gender in a study aiming at validating a measurement protocol and for evaluating the usability and acceptance level of an ICT equipment. The majority of the participants were men. A similar pattern is observed for Diabetes and nutrition, Gait and fall, Neurological diseases, Other, Rehabilitation and Stress and sleep. Women are only in majority for one of the groups in the Gait and fall study [30], and the Rehabilitation studies [41,42,48].

#### 3.4. Tests on Patients and Healthy Users

Information on whether the participants were patients and/or healthy was provided in 39/60 (65%) of the articles (Table 3). An additional four studies, [9,21,22,38] present the distribution of patients and/or healthy for some of the reported sub-studies. Two groups including 84 participants in total were representing patients and healthy participants in [19]. Seven articles [13,17,19,23,41,42,50] report on the conduction of studies with both patients and healthy. Two articles [9,22] contain results from several sub-studies and while not providing patient/healthy information for all sub-studies, claim to have used both patients and healthy participants during data collection. For several article categories, many of the studies reported information on both patients and healthy users.

Studying the articles from a health perspective, i.e., looking particularly at the article categories Asthma/COPD, Cardiovascular diseases, Diabetes and nutrition, Neurological diseases, and Stress and Sleep, the reporting and/or use of patients/healthy participants varies. Almost all participants in studies on Asthma/COPD and Neurological diseases were patients. Surprisingly, the Cardiovascular diseases [20,23] were conducted solely with healthy participants while another [21] and three of the sub-studies in [21,22] lack information on whether the participants were healthy or patients. Regarding Diabetes and Nutrition, two works [24,28] were conducted with patients, one study [27] was conducted with healthy participants while two articles [25,26] lack this information. Finally, regarding Stress and sleep, none of the studies report on studies with patients. Three articles [55,57,58] were conducted with healthy participants while the remaining three articles lack this information.

Studying the articles from a physical activity perspective, i.e., looking particularly at the article categories Gait and fall, Physical activity monitoring and Rehabilitation. No information on whether the participants were healthy or patients were provided in the articles falling under the Physical activity monitoring article category. None of the studies within Gait and fall used patients. The picture is mixed for the category Rehabilitation, two studies were conducted solely with patients [48,51] whereas [50] reports on two sub-studies conducted with patients and healthy participants respectively. One work [52] was conducted solely with healthy participants and two works [49,53] do not provide this information.

#### 4. Discussion and Conclusions

In this systematic review, we provide a qualitative synthesis on retrieved articles on using wearable body sensors for health monitoring. The articles found were categorized as relating to: Asthma/COPD, Cardiovascular diseases, Diabetes and Nutrition, Gait and fall, Neurological diseases, Physical activity recognition, Rehabilitation, Stress and sleep, and Additional. Section 3 provided a qualitative synthesis of the studies with respect to research methodology and participant demography, i.e., number of participants, age, gender and the distribution of healthy participants and patients. Using this information, we have identified a number of shortcomings. Below follows a discussion on these shortcomings in relation to prior research.

There are many age-related health issues such as changing biological factors, the onset of illnesses which are often chronic and the decline of cognitive abilities. For example, "fall prediction is a challenging problem due to the combination of intrinsic and extrinsic fall risk factors that contribute to a fall. Intrinsic factors include age, fall history, mobility impairments, sleep disturbances, and neurological disorders", pp. 1 [76]. It is reported in [77] that 35% of non-institutionalized adults had abnormal gait and that sleep disturbances are very common among older people. Further, chronic conditions affect physical activity levels, and activities such as rising from a chair is demanding

for older people [77]. It is clear that the whole motion pattern changes with age and the onset of illnesses related to the human locomotor system. Yet, the majority of the studies focusing on gait and fall in this review were simulations that include none or few old participants. This shortcoming is also discussed in [76], "It is evident that existing systems have mainly been tested in laboratory environments with controlled conditions and do not include frequent fallers and aging adults as test subjects.[..] future work should focus on longitudinal studies of fall detection and prediction systems in real-life conditions on a diverse group that includes frequent fallers, aging adults, and persons with neurological disorders." p.8 [76]. Not studying the sensor systems in real-life conditions affect the validity of the results since the performance is not studied in realistic conditions. The low number of studies with older people is also a shortcoming since age-related issues are not taken into consideration to a sufficient degree.

There are many differences between the two genders. As a first example, we want to mention the American Heart Association's (AHA) scientific statement from 2016 [78] on acute myocardial infarction (AMI) in women. "Sex differences occur in the pathophysiology and clinical presentation of MI and affect treatment delays.", p. 932 [78]. Further, AHA reports that the same perfusion therapies are recommended despite the fact that the risk of bleeding or other complications is higher among women. Further, women are being under-treated with guideline recommendations. This results in increased readmission, re-infarction, and death rates during the first year after a myocardial infarction. Cardiac rehabilitation is also underused and under-prescribed among women [78]. On the same lines, the results of a cohort study [79] with almost 5000 patients  $\mu_{age} > 65$  who were admitted to 366 US hospitals in the period 2003–2009, has found that women are less likely to receive optimal care at discharge. Yet, only two of the studies retrieved within the category Cardiovascular diseases provide information on the participants' gender. This is not the only shortcoming for studies on Cardiovascular diseases however. Several studies, or sub-studies, were conducted with very large age spans without the provision of a mean age. Others were conducted with young people or lacks information on age. Further, several works report on studies with healthy participants.

Hence, studies taking both genders into consideration, but also the age factor, are highly desired in the category Cardiovascular diseases. Not including information on gender and/or not considering gender/sex during data collection is a shortcoming regardless of the category to which a study belongs. It is argued in [80] that there are areas were specific data on women is lacking while specific data on men is missing in other areas.

Regitz-Zagrosek [80] outlines a number of differences between men and women. These include: women more frequently having anemia, women suffering from coronary artery disease in average ten years later than men, a higher frequency of boys having asthma in young ages while the frequency changes to young adulthood, diabetes increasing the risk for coronary heart disease more among women, and osteoporosis being more frequent in women but under-diagnosed in men. Osteoporosis disease is characterized by a decreased bone mass density and a disrupted normal trabecular architecture reducing bone strength [81]. Therefore, Osteoporosis increases the risk of fractures after a fall but no symptoms of the disease are shown until a fracture occurs [80]. According to [81], there are several factors relating to Osteoporosis which increases the risk of falling. These include the fear of falling, which increases the risk of falling [82,83]. In addition, [81] reports on studies discussing women with osteoporosis or low bone mass where fear of falling is associated with more falls [84], and the confidence in balance is related to balance and mobility [85]. Further, [84] reports that an increased thoracic kyphosis is associated with recent falls among women with Osteopororosis. I.e., women with thoracic kyphosis were more likely to have had a recent fall. Thoracic kyphosis is an abnormal convex curvature of the spine at chest height which is much more common among older women than men due to estrogen losses [86]. All these works [81–85] date from 2004-2011, hence it is astonishing that some articles retrieved within the article category Gait and fall have not reported information on gender and that some other articles were conducted solely with men. Hence, we argue that future studies in the

categories discussed in this article must take gender into consideration. This shortcoming was also highlighted in [2].

Undoubtedly, healthy participants and patients differ in many aspects. Yet, only 65% of the studies overall reported this information. A positive example here is the fact that the studies reported upon in the category Asthma/COPD were conducted almost entirely with patients. This indicates that the results in this area are reliable. On the contrary, none of the studies within Gait and fall, or Stress and sleep have reported that the studies were conducted with patients. Also [76,77] have previously discussed the shortcoming of not conducting studies with patients in the category Gait and fall. Considering the research question for this review article, we question the fact that 35% of the retrieved articles lack information on whether the participants were healthy or patients. We argue that the use of healthy participants, or not providing this information, affect the validity of the study results. Future studies need to consider the inclusion of patients to a further extent.

Studying the sample size in the reported studies, 56% of the articles report on studies conducted with up to 20 participants, and only 20% of the articles report on studies conducted with 51 or more participants. The distribution of numbers vary between categories. The majority of the studies reported in the categories Asthma/COPD, Gait and fall, Physical activity recognition, Rehabilitation, and Stress and sleep were conducted with up to 20 participants. We find the overall low number of participants a shortcoming and recommend that future studies are conducted with larger study samples. However, taking demographic factors, i.e., age, gender and healthy/patient into consideration is highly needed prior to increasing the sample sizes in studies on health monitoring using wearable body sensors.

**Author Contributions:** Search phrase, inclusion and exclusion criteria, A.K. and M.L.; literature search, A.K.; initial screening of articles, A.K.; selection of articles, A.K. and M.L.; article categorization, A.K.; tables and figures, A.K.; analysis of material in tables and figures, A.K. and M.L.; original draft preparation, A.K.; review and editing, A.K. and M.L. All authors have read and agree to the published version of the manuscript.

**Funding:** This research was conducted within the scope of the ESS-H+ (Embedded Sensor Systems for Health Plus). The project is funded by the Swedish Knowledge Foundation (project number: 20180158).

**Conflicts of Interest:** The authors declare no conflict of interest. The funder had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

#### References

- Moher, D.; Liberati, A.; Tetzlaff, J.; Altman, D.G.; the PRISMA Group. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *Ann. Internal Med.* 2009, 151, 264–269. [CrossRef]
- 2. Kekade, S.; Hseieh, C.H.; Islam, M.M.; Atique, S.; Mohammed Khalfan, A.; Li, Y.C.; Abdul, S.S. The usefulness and actual use of wearable devices among the elderly population. *Comput. Methods Programs Biomed.* **2018**, 153, 137–159. [CrossRef]
- 3. Li, Y.; Li, S.; Song, H.; Shao, B.; Yang, X.; Deng, N. Noninvasive blood pressure estimation with peak delay of different pulse waves. *Int. J. Distrib. Sens. Netw.* **2019**, *15*. [CrossRef]
- 4. Wu, X.; Wang, Y.; Chien, C.; Pottie, G. Self-calibration of sensor misplacement based on motion signatures. In Proceedings of the 2013 IEEE International Conference on Body Sensor Networks, Cambridge, MA, USA, 6–9 May 2013; pp. 1–5. [CrossRef]
- 5. Castro, D.; Coral, W.; Rodriguez, C.; Cabra, J.; Colorado, J. Wearable-Based Human Activity Recognition Using an IoT Approach. *J. Sens. Actuator Netw.* **2017**, *6*, 28. [CrossRef]
- 6. Rodriguez, C.; Castro, D.M.; Coral, W.; Cabra, J.L.; Velasquez, N.; Colorado, J.; Mendez, D.; Trujillo, L.C.; ACM. IoT system for Human Activity Recognition using BioHarness 3 and Smartphone. In Proceedings of the International Conference on Future Networks and Distributed Systems, Cambridge, UK, 19–20 July 2017. [CrossRef]

7. Doron, M.; Bastian, T.; Maire, A.; Dugas, J.; Perrin, E.; Gris, F.; Guillemaud, R.; Deschamps, T.; Bianchi, P.; Caritu, Y.; et al. Estimation of physical activity monitored during the day-to-day life by an autonomous wearable device (SVELTE project). In Proceedings of the 2013 35th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), Osaka, Japan, 3–7 July 2013; pp. 4629–4632. [CrossRef]

- 8. Xu, J.Y.; Chang, H.I.; Chien, C.; Kaiser, W.J.; Pottie, G.J. Context-driven, prescription-based personal activity classification: Methodology, architecture, and end-to-end implementation. *IEEE J. Biomed. Health Inform.* **2014**, *18*, 1015–1025. [CrossRef]
- 9. Xu, J.Y.; Wang, Y.; Barrett, M.; Dobkin, B.; Pottie, G.J.; Kaiser, W.J. Personalized multilayer maily life profiling through context enabled activity classification and motion reconstruction: An integrated system approach. *IEEE J. Biomed. Health Inform.* **2016**, *20*, 177–188. [CrossRef] [PubMed]
- 10. Velicu, O.R.; Madrid, N.M.; Seepold, R.; IEEE. Experimental sleep phases monitoring. In Proceedings of the 2016 3rd IEEE Embs International Conference on Biomedical and Health Informatics, Las Vegas, NV, USA, 24–27 February 2016; pp. 625–628.
- 11. Seeger, C.; Buchmann, A.; Van Laerhoven, K. An Event-based BSN Middleware That Supports Seamless Switching Between Sensor Configurations. In Proceedings of the 2nd ACM SIGHIT International Health Informatics Symposium, Miami, Fl, USA, 28–30 January 2012; pp. 503–512. [CrossRef]
- 12. Wannenburg, J.; Malekian, R. Body Sensor Network for Mobile Health Monitoring, a Diagnosis and Anticipating System. *IEEE Sens. J.* **2015**, *15*, 6839–6852. [CrossRef]
- 13. Bonnevie, T.; Gravier, F.E.; Elkins, M.; Dupuis, J.; Prieur, G.; Combret, Y.; Viacroze, C.; Debeaumont, D.; Robleda-Quesada, A.; Quieffin, J.; et al. People undertaking pulmonary rehabilitation are willing and able to provide accurate data via a remote pulse oximetry system: a multicentre observational study. *J. Physiother.* **2019**, *65*, 28–36. [CrossRef] [PubMed]
- Caulfield, B.; Kaljo, I.; Donnelly, S. Use of a consumer market activity monitoring and feedback device improves exercise capacity and activity levels in COPD. In Proceedings of the 2014 36th Annual International Conference of the IEEE Engineering in Medicine and Biology Society, Chicago, IL, USA, 26–30 August 2014; pp. 1765–1768. [CrossRef]
- 15. Estrada, L.; Torres, A.; Sarlabous, L.; Jané, R. Evaluating respiratory muscle activity using a wireless sensor platform. In Proceedings of the 2016 38th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), Orlando, FL, USA, 16–20 August 2016; pp. 5769–5772. [CrossRef]
- 16. Katsaras, T.; Milsis, A.; Rizikari, M.; Saoulis, N.; Varoutaki, E.; Vontetsianos, A. The use of the "Healthwear" wearable system in chronic patients' early hospital discharge: Control randomized clinical trial. In Proceedings of the 2011 5th International Symposium on Medical Information and Communication Technology, Montreux, Switzerland, 27–30 March 2011; pp. 143–146. [CrossRef]
- 17. Naranjo-Hernández, D.; Talaminos-Barroso, A.; Reina-Tosina, J.; Roa, L.M.; Barbarov-Rostan, G.; Cejudo-Ramos, P.; Márquez-Martín, E.; Ortega-Ruiz, F. Smart vest for respiratory rate monitoring of copd patients based on non-contact capacitive sensing. *Sensors* 2018, 18, 2144. [CrossRef]
- 18. Huang, A.; Xu, W.; Li, Z.; Xie, L.; Sarrafzadeh, M.; Li, X.; Cong, J. System Light-Loading Technology for mHealth: Manifold-Learning-Based Medical Data Cleansing and Clinical Trials in WE-CARE Project. *IEEE J. Biomed. Health Inform.* **2014**, *18*, 1581–1589. [CrossRef] [PubMed]
- 19. Huang, A.; Chen, C.; Bian, K.; Duan, X.; Chen, M.; Gao, H.; Meng, C.; Zheng, Q.; Zhang, Y.; Jiao, B.; et al. WE-CARE: An intelligent mobile telecardiology system to enable mHealth applications. *IEEE J. Biomed. Health Inform.* **2014**, *18*, 693–702. [CrossRef] [PubMed]
- Javaid, A.Q.; Chang, I.S.; Mihailidis, A. Ballistocardiogram Based Identity Recognition: Towards Zero-Effort Health Monitoring in an Internet-of-Things (IoT) Environment. In Proceedings of the 2018 40th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), Honolulu, HI, USA, 18–21 July 2018, Volume 2018; pp. 3326–3329. [CrossRef]
- 21. Raad, M.W.; Sheltami, T.; Deriche, M. A Ubiquitous Telehealth System for the Elderly. In *Internet of Things: User-Centric Iot, Pt I*; Lecture Notes of the Institute for Computer Sciences Social Informatics and Telecommunications Engineering; Giaffreda, R., Vieriu, R.L., Pasher, E., Bendersky, G., Jara, A.J., Rodrigues, J., Dekel, E., Mandler, B., Eds.; Springer: Berlin, Germany, 2015; Volume 150, pp. 159–166. [CrossRef]
- 22. Simjanoska, M.; Gjoreski, M.; Gams, M.; Bogdanova, A.M. Non-invasive blood pressure estimation from ECG using machine learning techniques. *Sensors* **2018**, *18*, 1160. [CrossRef] [PubMed]

23. Susič, T.P.; Stanič, U. Penetration of the ICT technology to the health care primary sector—Ljubljana PILOT. In Proceedings of the 2016 39th International Convention on Information and Communication Technology, Electronics and Microelectronics (MIPRO), Opatija, Croatia, 30 May–3 June 2016; pp. 436–441. [CrossRef]

- 24. Al-Taee, M.A.; Al-Nuaimy, W.; Al-Ataby, A.; Muhsin, Z.J.; Abood, S.N.; IEEE. Mobile Health Platform for Diabetes Management Based on the Internet-of-Things. In Proceedings of the 2015 IEEE Jordan Conference on Applied Electrical Engineering and Computing Technologies, New York, NY, USA, 3–5 November 2015. [CrossRef]
- 25. Alshurafa, N.; Kalantarian, H.; Pourhomayoun, M.; Sarin, S.; Liu, J.J.; Sarrafzadeh, M. Non-invasive monitoring of eating behavior using spectrogram analysis in a wearable necklace. In Proceedings of the 2014 IEEE Healthcare Innovation Conference (HIC), Seattle, WA, USA, 8–10 October 2014; pp. 71–74. [CrossRef]
- 26. Alshurafa, N.; Kalantarian, H.; Pourhomayoun, M.; Liu, J.J.; Sarin, S.; Shahbazi, B.; Sarrafzadeh, M. Recognition of Nutrition Intake Using Time-Frequency Decomposition in a Wearable Necklace Using a Piezoelectric Sensor. *IEEE Sens. J.* 2015, 15, 3909–3916. [CrossRef]
- 27. Dong, B.; Biswas, S. Meal-time and duration monitoring using wearable sensors. *Biomed. Signal Process. Control* **2017**, 32, 97–109. [CrossRef]
- 28. Onoue, T.; Goto, M.; Kobayashi, T.; Tominaga, T.; Ando, M.; Honda, H.; Yoshida, Y.; Tosaki, T.; Yokoi, H.; Kato, S.; et al. Randomized controlled trial for assessment of Internet of Things system to guide intensive glucose control in diabetes outpatients: Nagoya Health Navigator Study protocol. *Nagoya J. Med. Sci.* 2017, 79, 323–329. [CrossRef] [PubMed]
- 29. Atallah, L.; Wiik, A.; Jones, G.G.; Lo, B.; Cobb, J.P.; Amis, A.; Yang, G.Z. Validation of an ear-worn sensor for gait monitoring using a force-plate instrumented treadmill. *Gait Posture* **2012**, *35*, 674–676. [CrossRef]
- 30. Godfrey, A.; Din, S.D.; Barry, G.; Mathers, J.C.; Rochester, L. Within trial validation and reliability of a single tri-axial accelerometer for gait assessment. In Proceedings of the 2014 36th Annual International Conference of the IEEE Engineering in Medicine and Biology Society, Chicago, IL, USA, 26–30 August 2014; pp. 5892–5895. [CrossRef]
- 31. Lee, J.K.; Robinovitch, S.N.; Park, E.J. Inertial Sensing-Based Pre-Impact Detection of Falls Involving Near-Fall Scenarios. *IEEE Trans. Neural Syst. Rehabil. Eng.* **2015**, 23, 258–266. [CrossRef]
- 32. Liang, D.; Zhao, G.; Guo, Y.; Wang, L. Pre-impact & impact detection of falls using wireless Body Sensor Network. In Proceedings of the 2012 IEEE-EMBS International Conference on Biomedical and Health Informatics, Hong Kong, China, 5–7 January 2012; pp. 763–766. [CrossRef]
- 33. Liang, S.; Chu, T.; Lin, D.; Ning, Y.; Li, H.; Zhao, G. Pre-impact Alarm System for Fall Detection Using MEMS Sensors and HMM-based SVM Classifier. In Proceedings of the 2018 40th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), Honolulu, HI, USA, 18–21 July 2018; pp. 4401–4405. [CrossRef]
- 34. Paiman, C.; Lemus, D.; Short, D.; Vallery, H. Observing the State of Balance with a Single Upper-Body Sensor. *Front. Robot. AI* **2016**, *3.* [CrossRef]
- 35. Tino, A.; Carvalho, M.; Preto, N.F.; McConville, K.M.V. Wireless vibrotactile feedback system for postural response improvement. In Proceedings of the 2011 33rd Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), Boston, MA, USA, 30 August–3 September 2011; Volume 2011, pp. 5203–2506. [CrossRef]
- 36. Williams, B.; Allen, B.; True, H.; Fell, N.; Levine, D.; Sartipi, M.; IEEE. A Real-time, Mobile Timed Up and Go System. In Proceedings of the 2015 IEEE 12th International Conference on Wearable and Implantable Body Sensor Networks (BSN), Cambridge, MA, USA, 9–12 June 2015. [CrossRef]
- 37. Wu, Y.; Su, Y.; Feng, R.; Yu, N.; Zang, X. Wearable-sensor-based pre-impact fall detection system with a hierarchical classifier. *Measurement* **2019**, *140*, 283–292. [CrossRef]
- 38. Zhao, G.; Mei, Z.; Liang, D.; Kamen, I.; Guo, Y.; Wang, Y.; Wang, L. Exploration and Implementation of a Pre-Impact Fall Recognition Method Based on an Inertial Body Sensor Network. *Sensors* **2012**, 12, 15338–15355. [CrossRef]
- 39. Zhong, R.; Rau, P.L.P.; Yan, X. Gait Assessment of Younger and Older Adults with Portable Motion-Sensing Methods: A User Study. *Mob. Inf. Syst.* **2019**, 2019. [CrossRef]
- 40. Giuberti, M.; Ferrari, G.; Contin, L.; Cimolin, V.; Azzaro, C.; Albani, G.; Mauro, A. Automatic UPDRS Evaluation in the Sit-to-Stand Task of Parkinsonians: Kinematic Analysis and Comparative Outlook on the Leg Agility Task. *IEEE J. Biomed. Health Inform.* **2015**, *19*, 803–814. [CrossRef] [PubMed]

41. Gong, J.; Lach, J.; Qi, Y.; Goldman, M.D. Causal analysis of inertial body sensors for enhancing gait assessment separability towards multiple sclerosis diagnosis. In Proceedings of the 2015 IEEE 12th International Conference on Wearable and Implantable Body Sensor Networks (BSN), Cambridge, MA, USA, 9–12 June 2015. [CrossRef]

- 42. Gong, J.; Qi, Y.; Goldman, M.D.; Lach, J. Causality Analysis of Inertial Body Sensors for Multiple Sclerosis Diagnostic Enhancement. *IEEE J. Biomed. Health Inform.* **2016**, *20*, 1273–1280. [CrossRef] [PubMed]
- 43. Kuusik, A.; Alam, M.M.; Kask, T.; Gross-Paju, K. Wearable m-assessment system for neurological disease patients. In Proceedings of the 2018 IEEE 4th World Forum on Internet of Things (WF-IoT), Singapore, Singapore, 5–8 February 2018; pp. 201–206. [CrossRef]
- 44. Sok, P.; Xiao, T.; Azeze, Y.; Jayaraman, A.; Albert, M.V. Activity recognition for incomplete spinal cord injury subjects using hidden markov models. *IEEE Sens. J.* **2018**, *18*, 6369–6374. [CrossRef]
- 45. Stamate, C.; Magoulas, G.D.; Kueppers, S.; Nomikou, E.; Daskalopoulos, I.; Luchini, M.U.; Moussouri, T.; Roussos, G. Deep learning Parkinson's from smartphone data. In Proceedings of the 2017 IEEE International Conference on Pervasive Computing and Communications, PerCom 2017, Kona, HI, USA, 13–17 March 2017; pp. 31–40. [CrossRef]
- 46. Stamate, C.; Magoulas, G.D.; Kueppers, S.; Nomikou, E.; Daskalopoulos, I.; Jha, A.; Pons, J.S.; Rothwell, J.; Luchini, M.U.; Moussouri, T.; et al. The cloudUPDRS app: A medical device for the clinical assessment of Parkinson's Disease. *Pervasive Mob. Comput.* **2018**, *43*, 146–166. [CrossRef]
- 47. Rednic, R.; Gaura, E.; Brusey, J.; Kemp, J. Wearable posture recognition systems: Factors affecting performance. In Proceedings of the 2012 IEEE-EMBS International Conference on Biomedical and Health Informatics, Hong Kong, China, 5–7 January 2012; pp. 200–203. [CrossRef]
- 48. Argent, R.; Slevin, P.; Bevilacqua, A.; Neligan, M.; Daly, A.; Caulfield, B. Wearable sensor-based exercise biofeedback for orthopaedic rehabilitation: A mixed methods user evaluation of a prototype system. *Sensors* **2019**, *19*, 432. [CrossRef]
- 49. Banos, O.; Moral-Munoz, J.A.; Diaz-Reyes, I.; Arroyo-Morales, M.; Damas, M.; Herrera-Viedma, E.; Hong, C.S.; Lee, S.; Pomares, H.; Rojas, I.; et al. MDurance: A novel mobile health system to support trunk endurance assessment. *Sensors* 2015, *15*, 13159–13183. [CrossRef]
- 50. Lee, S.I.; Adans-Dester, C.P.; Grimaldi, M.; Dowling, A.V.; Horak, P.C.; Black-Schaffer, R.M.; Bonato, P.; Gwin, J.T. Enabling stroke rehabilitation in home and community settings: A wearable sensor-based approach for upper-limb motor training. *IEEE J. Transl. Eng. Health Med.* **2018**, *6*. [CrossRef]
- 51. Timmermans, A.A.A.; Seelen, H.A.M.; Geers, R.P.J.; Saini, P.K.; Winter, S.; te Vrugt, J.; Kingma, H. Sensor-Based Arm Skill Training in Chronic Stroke Patients: Results on Treatment Outcome, Patient Motivation, and System Usability. *IEEE Trans. Neural Syst. Rehabil. Eng.* 2010, 18, 284–292. [CrossRef]
- 52. Whelan, D.F.; O'Reilly, M.A.; Ward, T.E.; Delahunt, E.; Caulfield, B. Technology in rehabilitation: Comparing personalised and global classification methodologies in evaluating the squat exercise with wearable IMUs. *Methods Inf. Med.* **2017**, *56*, 361–369. [CrossRef]
- 53. Xu, J.K.; Lee, U.H.; Bao, T.; Huang, Y.J.; Sienko, K.H.; Shull, P.B.; IEEE. Wearable sensing and haptic feedback research platform for gait retraining. In Proceedings of the 2017 IEEE 14th International Conference on Wearable and Implantable Body Sensor Networks, Eindhoven, The Netherlands, 9–12 May 2017; pp. 125–128.
- 54. Lin, C.; Gamble, J.; Yang, Y.; Wang, J. Estimating the influence of chronotype and social zeitgebers on circadian rhythms using an accelerometer-based sensor network. In Proceedings of the 2012 IEEE-EMBS International Conference on Biomedical and Health Informatics, Hong Kong, China, 5–7 January 2012; pp. 285–288. [CrossRef]
- 55. Nakamura, T.; Goverdovsky, V.; Morrell, M.J.; Mandic, D.P. Automatic Sleep Monitoring Using Ear-EEG. *IEEE J. Transl. Eng. Health Med.* **2017**, *5*, 1–8. [CrossRef] [PubMed]
- 56. Parnandi, A.; Gutierrez-Osuna, R. Physiological Modalities for Relaxation Skill Transfer in Biofeedback Games. *IEEE J. Biomed. Health Inform.* **2017**, 21, 361–371. [CrossRef] [PubMed]
- 57. Uday, S.; Jyotsna, C.; Amudha, J.; IEEE. Detection of Stress using Wearable Sensors in IoT Platform. In Proceedings of the 2018 Second International Conference on Inventive Communication and Computational Technologies (ICICCT), Coimbatore, India, 20–21 April 2018; pp. 492–498. [CrossRef]

Sensors **2020**, 20, 1502 20 of 21

58. Umemura, G.S.; Pinho, J.P.; Furtado, F.; Gonçalves, B.S.B.; Fomer-Cordero, A. Comparison of sleep parameters assessed by actigraphy of healthy young adults from a small town and a megalopolis in an emerging country. In Proceedings of the 2017 IEEE Healthcare Innovations and Point of Care Technologies (HI-POCT), Bethesda, MD, USA, 6–8 November 2017; pp. 200–203. [CrossRef]

- 59. Ayzenberg, Y.; Picard, R.W. FEEL: A System for Frequent Event and Electrodermal Activity Labeling. *IEEE J. Biomed. Health Inform.* **2014**, *18*, 266–277. [CrossRef] [PubMed]
- Pagán, J.; Risco-Martín, J.L.; Moya, J.M.; Ayala, J.L. Grammatical Evolutionary Techniques for Prompt Migraine Prediction. In Proceedings of the Genetic and Evolutionary Computation Conference Denver, CO, USA, 20–24 July 2016; pp. 973–980. [CrossRef]
- 61. Rawashdeh, M.; Al-Qurishi, M.; Al-Rakhami, M.; Al-Quraishi, M.S. A multimedia cloud-based framework for constant monitoring on obese patients. In Proceedings of the 2017 IEEE International Conference on Multimedia & Expo Workshops (ICMEW), Hong Kong, China, 10–14 July 2017; pp. 139–144. [CrossRef]
- 62. Wu, W.; Pirbhulal, S.; Sangaiah, A.K.; Mukhopadhyay, S.C.; Li, G. Optimization of signal quality over comfortability of textile electrodes for ECG monitoring in fog computing based medical applications. *Future Gener. Comput. Syst.* **2018**, *86*, 515–526. [CrossRef]
- 63. Buonocore, C.M.; Rocchio, R.A.; Roman, A.; King, C.E.; Sarrafzadeh, M. Wireless Sensor-Dependent Ecological Momentary Assessment for Pediatric Asthma mHealth Applications. In Proceedings of the Second IEEE/ACM International Conference on Connected Health: Applications, Systems and Engineering Technologies, Philadelphia, PA, USA, 17–19 July 2017; pp. 137–146. [CrossRef]
- 64. Depolli, M.; Avbelj, V.; Trobec, R.; Kališnik, J.M.; Tadej, K.; Susič, A.P.; Stanič, U.; Semeja, A. PCARD platform for mhealth monitoring. *Informatica* **2016**, *40*, 117–123.
- 65. Ghazal, M.; Khalil, Y.A.; Dehbozorgi, F.J.; Alhalabi, M.T. An integrated caregiver-focused mHealth framework for elderly care. In Proceedings of the 11th IEEE International Conference on Wireless and Mobile Computing, Networking and Communications, WiMob 2015, Abu Dhabi, UAE, 19–21 October 2015; pp. 238–245. [CrossRef]
- 66. Anupama, K.R.; Adarsh, R.; Pahwa, P.; Ramachandran, A. Machine Learning-Based Techniques for Fall Detection in Geriatric Healthcare Systems. In Proceedings of the 2018 9th International Conference on Information Technology in Medicine and Education (ITME), Hangzhou, China, 19–21 October 2018; pp. 232–237. [CrossRef]
- 67. Boutellaa, E.; Kerdjidj, O.; Ghanem, K. Covariance matrix based fall detection from multiple wearable sensors. *J. Biomed. Inform.* **2019**, 103189. [CrossRef]
- 68. Memedi, M.; Tshering, G.; Fogelberg, M.; Jusufi, I.; Kolkowska, E.; Klein, G. An Interface for IoT: Feeding Back Health-Related Data to Parkinson's Disease Patients. *J. Sens. Actuator Netw.* **2018**, 7. [CrossRef]
- 69. Dobbins, C.; Rawassizadeh, R.; Momeni, E. Detecting physical activity within lifelogs towards preventing obesity and aiding ambient assisted living. *Neurocomputing* **2017**, 230, 110–132. [CrossRef]
- 70. Argent, R.; Slevin, P.; Bevilacqua, A.; Neligan, M.; Daly, A.; Caulfield, B. Clinician perceptions of a prototype wearable exercise biofeedback system for orthopaedic rehabilitation: A qualitative exploration. *BMJ Open* **2018**, *8*. [CrossRef]
- 71. Zhuang, Y.; Song, C.; Wang, A.; Lin, F.; Li, Y.; Gu, C.; Li, C.; Xu, W. SleepSense: Non-invasive sleep event recognition using an electromagnetic probe. In Proceedings of the 2015 IEEE 12th International Conference on Wearable and Implantable Body Sensor Networks (BSN), Cambridge, MA, USA, 9–12 June 2015. [CrossRef]
- 72. Amira, T.; Dan, I.; Az-eddine, B.; Ngo, H.H.; Said, G.; Katarzyna, W. Monitoring chronic disease at home using connected devices. In Proceedings of the 2018 13th Annual Conference on System of Systems Engineering (SoSE), Paris, France, 19–22 June 2018; pp. 400–407. [CrossRef]
- 73. Cortinas, R.; Gonzaga, J.M.; Green, A.R.; Saulenas, A.M.; BuSha, B.F. TCNJ Athlete Tracker. In Proceedings of the 2015 41st Annual Northeast Biomedical Engineering Conference (NEBEC), Troy, NY, USA, 17–19 April 2015. [CrossRef]
- 74. Hörmann, T.; Hesse, M.; Adams, M.; Rückert, U. A Software Assistant for User-Centric Calibration of a Wireless Body Sensor. In Proceedings of the 2016 IEEE 13th International Conference on Wearable and Implantable Body Sensor Networks (BSN), San Francisco, CA, USA, 14–17 June 2016; pp. 183–188. [CrossRef]
- 75. Warmerdam, L.; Riper, H.; Klein, M.; van den Ven, P.; Rocha, A.; Ricardo Henriques, M.; Tousset, E.; Silva, H.; Andersson, G.; Cuijpers, P. Innovative ICT solutions to improve treatment outcomes for depression: the ICT4Depression project. *Stud. Health Technol. Inform.* **2012**, *181*, 339–343.

Sensors **2020**, 20, 1502 21 of 21

76. Rajagopalan, R.; Litvan, I.; Jung, T.P. Fall Prediction and Prevention Systems: Recent Trends, Challenges, and Future Research Directions. *Sensors* **2017**, *17*, 2509. [CrossRef]

- 77. Tedesco, S.; Barton, J.; O'Flynn, B. A review of activity trackers for senior citizens: Research perspectives, commercial landscape and the role of the insurance industry. *Sensors* **2017**, *17*, 1277. [CrossRef]
- 78. Mehta, L.S.; Beckie, T.M.; DeVon, H.A.; Grines, C.L.; Krumholz, H.M.; Johnson, M.N.; Lindley, K.J.; Vaccarino, V.; Wang, T.Y.; Watson, K.E.; et al. Acute Myocardial Infarction in Women A Scientific Statement From the American Heart Association. *Circulation* 2016, 133, 916–947. [CrossRef] [PubMed]
- 79. Li, S.; Fonarow, G.C.; Mukamal, K.J.; Liang, L.; Schulte, P.J.; Smith, E.E.; DeVore, A.; Hernandez, A.F.; Peterson, E.D.; Bhatt, D.L. Sex and Race/Ethnicity–Related Disparities in Care and Outcomes After Hospitalization for Coronary Artery Disease Among Older Adults. *Circ. Cardiovasc. Qual. Outcomes* 2016, 9, 36–44. [CrossRef] [PubMed]
- 80. Regitz-Zagrosek, V. Sex and gender differences in health. Science & Society Series on Sex and Science. *EMBO Rep.* **2012**, *13*, 596–603. [CrossRef] [PubMed]
- 81. Smulders, E.; van Lankveld, W.; Laan, R.; Duysens, J.; Weerdesteyn, V. Does osteoporosis predispose falls? a study on obstacle avoidance and balance confidence. *BMC Musculoskelet*. *Disord*. **2011**, *12*, 1. [CrossRef]
- 82. Scheffer, A.; Schuurmans, M.; van Dijk, N.; van der Hooft, T.; de Rooij, S. Fear of falling: measurement strategy, prevalence, risk factors and consequences among older persons. *Age Ageing* **2008**, *37*, 19–24. [CrossRef]
- 83. Delbaere, K.; Crombez, G.; Vanderstraeten, G.; Willems, T.; Cambier, D. Fear-related avoidance of activities, falls and physical frailty. A prospective community-based cohort study. *Age Ageing* **2004**, *33*, 368–373. [CrossRef]
- 84. Arnold, D.; Busch, A.; Schachter, C.; Harrison, L.; Olsynski, W. The relationship of intrinsic fall risk factors to a recent history of falling in older women with osteoporosis. *J. Orthop. Sports Phys. Ther.* **2005**, *35*, 452–460. [CrossRef]
- 85. Liu-Ambrose, T.; Khan, K.; Donaldson, M.; Eng, J.; Lord, S.; McKay, H. Falls-related self-efficacy is independently associated with balance and mobility in older women with low bone mass. *J. Gerontol. Ser. A* **2006**, *51*, 832–838. [CrossRef]
- 86. A Patient's Guide to Adult Kyphosis. Available online: https://www.umms.org/ummc/health-services/orthopedics/services/spine/patient-guides/adult-kyphosis (accessed on 16 January 2020).



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).